

Light adjoint scalars and unification at the string scale.

Mar Bastero-Gil^a and Biswajoy Brahmachari^b

(a) Scuola Internazionale Superiore di Studi Avanzati
34013 Trieste, ITALY.

(b) International Centre For Theoretical Physics,
34100 Trieste, ITALY.

Abstract

Following the suggestion of Bachas, Fabre and Yanagida (BFY), we analyze the gauge coupling unification at the two-loop order, in a supersymmetric scenario where scalars belonging to the adjoint representations contribute to the evolution of the couplings from intermediate scales onward, and the unification scale is pushed towards the string scale. Thereafter, we compare the masses of these adjoint scalars to the scale at which the hidden sector gauge coupling reaches the non-perturbative limit at the two-loop order for various possible hidden sector gauge groups motivated by the conjecture of BFY that the masses of these adjoint scalars are related to gaugino condensation. We also compute the predictions for the top and bottom quark masses in this scenario and compare them with those of MSSM. The predicted bottom mass improves in the BFY scenario for smaller values of α_s .

The gauge couplings unify at $M_{string} = 2 \times 10^{16} GeV$ when Minimal Supersymmetric Standard Model (MSSM) degrees of freedom contributes to the evolution of the gauge couplings. However, if the field theory of MSSM is a low energy remnant of the string theory, there exists a noteworthy discrepancy. This field theoretic unification scale is different from the scale of the string theory, M_{string} , which is fixed by the intrinsic scale of the string theory, namely M_{Plank} by the relation,

$$M_{string} = g_{string} M_{Plank}. \quad (1)$$

One loop string effects could lower this tree level value of the string scale somewhat, and one calculates [1, 2] that the string unification scale is modified to,

$$M_{string} = g_{string} \times 5.27 \times 10^{17} \text{ GeV} \simeq 5.27 \times 10^{17} \text{ GeV}, \quad (2)$$

where we have assumed $g_{string} = O(1)$. Consequently, the scale M_{string} is higher than the scale M_{GUT} by a factor of 20 approximately.

In the literature one finds several approaches to rectify the aforementioned difference between the string scale and the unification scale. There can be intermediate symmetry groups [3, 4], which alter the running of the gauge couplings and likewise the unification occurs at the scale of M_{string} . Non-standard hypercharge normalizations, which arise in various string models, can also accommodate the apparent mismatch between the string scale and the unification scale [5]. The heavy string threshold corrections from the string states [6] at the Plank scale or below have been used to reconcile the mismatch between the string scale and the unification scale. Non-standard exotic matter [7] has also been shown to serve the purpose.

The string models having a $G \times G$ structure, when broken to the diagonal subgroup, naturally contains adjoint scalars with zero hypercharge. Bachas, Fabre and Yanagida (BFY) have pointed out [9] that, if the mass of these zero-hypercharge adjoint scalars lie in the well motivated intermediate scale $M_I \sim M^{2/3} m_{susy}^{1/3} \sim 10^{13} \text{ GeV}$, the unification scale is pushed up to the string scale at the one-loop level. In this paper we present a two-loop analysis of this scenario and find ranges of the masses of the $SU(2)$ and $SU(3)$ adjoint scalars, namely, M_2 and M_3 for the allowed

range of $\alpha_s(m_Z)$ [8] which gives rise to the gauge coupling unification at the scale M_{string} . At the two-loop level there is also a mild variation of the unification scale to the ratio $\tan \beta = \langle H_2 \rangle / \langle H_1 \rangle$, because at the two-loop level, the running of the gauge couplings are affected by the simultaneous running of the Yukawa couplings. The adjoint scalars have no Yukawa couplings with the ordinary Fermions and so we consider the effect of only the ordinary Yukawa couplings on the evolution of gauge couplings.

At the two-loop level the evolution of the gauge couplings are governed by the equation,

$$\frac{d\alpha_i}{dt} = \frac{b_i}{2\pi}\alpha_i^2 + \sum_j \frac{b_{ij}}{8\pi^2}\alpha_i^2\alpha_j - \sum_k \frac{a_{ik}}{8\pi^2}\alpha_i^2 Y_k, \quad (3)$$

where,

$$b_i = \begin{pmatrix} 2n_f + \frac{3}{5}n_d \\ -6 + 2n_f + n_d + 2\Theta(\mu - M_2) \\ -9 + 2n_f + 3\Theta(\mu - M_3) \end{pmatrix} \quad a_{ik} = \begin{pmatrix} \frac{26}{5} & \frac{14}{5} & \frac{18}{5} \\ 6 & 6 & 2 \\ 4 & 4 & 0 \end{pmatrix}, \quad (4)$$

and,

$$b_{ij} = \begin{pmatrix} \frac{38}{15}n_f + \frac{9}{25}n_d & \frac{6}{5}n_f + \frac{9}{5}n_d & \frac{88}{15}n_f \\ \frac{2}{5}n_f + \frac{3}{5}n_d & -24 + 14n_f + 7n_d + 24\Theta(\mu - M_2) & 8n_f \\ \frac{11}{5}n_f & 3n_f & -54 + \frac{68}{3}n_f + 54\Theta(\mu - M_3) \end{pmatrix}, \quad (5)$$

where, $i, j=1, 3$; $k=t, b, \tau$; n_d are the number of Higgs doublets, n_f is the number of Fermion generations, $t = \ln \mu$ and $\Theta(x) = 1$, whenever $x \geq 0$. Fixing the scale $M_X = M_{string}$ we can find three quantities, from a system of three evolution equations, using as inputs the quantities $\alpha_1(m_Z)$, $\alpha_2(m_Z)$ and $\alpha_s(m_Z)$. The evaluated unknowns are α_G , M_2 and M_3 . The calculated values of M_2 and M_3 are plotted in Figure (1.b). The magnitude of M_3 has an appreciable sensitivity to the input value of $\alpha_s(m_Z)$. This is also displayed in the Figures (1.a) and (1.b).

It is noticeable from Figure (1.b) that the adjoint scalars having the mass near $10^{12} - 10^{14}$ GeV. Numerically, this scale is roughly near $M_{Plank}^{2/3} m_{susy}^{1/3}$. This is a striking coincidence, as noted by BFK. Now to check further, we will assume the unification of the gauge couplings of the hidden and observable sectors at the scale M_{string} . Using the initial value of the observable sector coupling $\alpha_G(M_{string})$ we evolve the hidden gauge coupling backwards and search for the scale M_C




Figure 1: The values of M_2 and M_3 for various values of the unification scale are in Figure (b). The shaded region refers to the variation of $\tan \beta$. In Figure (a) we have compared the values of M_2 and M_3 with the condensation scale M_C when M_{string} is 5.27×10^{17} GeV and G_H is taken as $SU(3)$, $SU(5)$ and $SO(10)$.

at which the hidden sector gauge coupling becomes non-perturbative. We have, for the simplicity, considered only matter-free gauge groups which are the subgroups of E_8 , eg, $SU(2)$, $SU(3)$, $SU(5)$ or $SO(10)$. In Figure (1.a) we have compared the scale M_C with M_2 and M_3 when G_H is $SU(3)$, $SU(5)$ and $SO(10)$ respectively. The condensation scale, when G_H is $SO(10)$ is large compared to the scales M_2 and M_3 . The choice $SU(3)$ as the hidden group returns a closer value of M_C to the scales M_3 and M_2 than if G_H is chosen as $SU(5)$. A comment about the results in two-loop calculation is in order. Consider the case when G_H is $SU(5)$ and $M_{string} = 2 \times 10^{17}$ GeV. In the one loop case the scale M_C is $10^{13.5}$ GeV but at the two-loop case this scale becomes $10^{15.2}$ GeV. Hence, the two-loop calculation shows a departure from the one-loop expectation and brings M_C closer to the intermediate scales M_2 and M_3 when the hidden group is $SU(3)$. This can be seen in Figure (2.a), where the running of the couplings are plotted. The running of the couplings in the MSSM are plotted in Figure (2.b) for comparison¹. We have taken $SU(5)$ as the unification

¹One of the authors B.B acknowledges discussions with Q. Shafi on the running of the hidden sector gauge coupling.




Figure 2: MSSM unification with a $SU(5)$ GUT with the minimal Higgs content is Figure (b). The $SU(5)$ coupling at the string scale is assumed to be unified with the hidden sector at the string scale. Figure (a) is string unification with adjoint moduli at the intermediate scale.

group from $M_X \simeq 2 \times 10^{16}$ GeV up to M_{string} .

Before we conclude, we give some results of a two-loop Yukawa coupling analysis of this model. The relevant beta function coefficients are summarized in Ref [4]. The Top quark mass is determined from the quasi-infrared fixed point of the top quark Yukawa coupling [11]. The presence of the adjoints alter the running of the gauge couplings above the intermediate scale which has secondary effects on the running of the Yukawa couplings. Even though we do not expect a large change in the individual Yukawa couplings at low energy, the ratios of the Yukawa couplings are sensitive to the presence of these extra matter. To check these issues, we have plotted the prediction of the top quark and the bottom quark masses in the Figures (3.a) and (3.b). The bottom quark mass can be calculated under the well-known assumption of b - τ unification². The prediction of the bottom quark mass, having been calculated from the *ratio* of the bottom and τ Yukawa coupling, differs notably when the extra adjoint matter is present. To see this we have

² b - τ unification is natural in GUT models like $SU(5)$ or $SO(10)$. Here, we consider a similar scenario in string unification.

G_H	$C_2(G)$	$M_{string} = 2 \times 10^{17}$	$M_{string} = 5.27 \times 10^{17}$	$M_{string} = 7 \times 10^{17}$
$SU(2)$	2	$M_C = 10^{10.1} (10^{7.8})$	$M_C = 10^{10.9} (10^{8.7})$	$M_C = 10^{11.1} (10^{9.0})$
$SU(3)$	3	$M_C = 10^{13.0} (10^{11.0})$	$M_C = 10^{13.7} (10^{11.7})$	$M_C = 10^{13.8} (10^{11.9})$
$SU(5)$	5	$M_C = 10^{15.2} (10^{13.5})$	$M_C = 10^{15.7} (10^{14.1})$	$M_C = 10^{15.9} (10^{14.3})$
$SO(10)$	8	$M_C = 10^{16.3} (10^{14.9})$	$M_C = 10^{16.7} (10^{15.5})$	$M_C = 10^{16.9} (10^{15.6})$
$\alpha_s = 0.117$	$SU(2)$ adjoint	$M_2 = 10^{14.8}$	$M_2 = 10^{14.1}$	$M_2 = 10^{13.9}$
$\alpha_s = 0.117$	$SU(3)$ adjoint	$M_3 = 10^{14.2}$	$M_3 = 10^{13.4}$	$M_3 = 10^{13.1}$

Table 1: The masses of the adjoint scalars of $SU(2)$ and $SU(3)$ that leads to gauge coupling unification at the scale M_{string} is in the two bottom lines of the table. The values quoted are for $\alpha_s = 0.117$. It can be compared to the scale M_C at which the hidden sector coupling becomes of order one for various gauge groups. For comparison the corresponding one-loop values are shown inside the brackets. The gauge groups have been designated by their respective quadratic casimirs.

compared the present case with that of MSSM in Figure (3.a) and (3.b) in dashed lines. The bands in the figures (3.a) and (3.b) are obtained by varying $\tan\beta$ in the range 5-60 and M_{string} in the range $2 - 7 \times 10^{17}$ GeV. In MSSM the unification occurs for only a narrow region in the α_s space, whereas, the presence of adjoint moduli opens up the parameter space further and as a welcome result unification occurs for a larger range in α_s . This fact is reflected in Figure (3) where the prediction of the top and bottom quark masses are displayed as narrow lines, whereas in the present case with adjoint moduli the predictions are wide bands. The bottom quark mass can be within the experimental range for lower values of α_s , for which the prediction in the MSSM case is unavailable due to the absence of gauge coupling unification and hence b - τ unification. In Table 2 we have compared the predictions of m_t^{pole} , m_b^{pole} and $\tan\beta$ in the MSSM with those obtained in the presence of adjoints for the required value of $\alpha_s(m_Z)$ which gives rise to unification in MSSM [Note that this specific value of α_s depends on Y_b or equivalently on $\tan\beta$]. Typically, the value of m_t^{pole} diminishes while m_b^{pole} increases slightly in the BFY scenario.

In conclusion, we have computed the two-loop running of the gauge couplings in a supersymmetric scenario with extra adjoint matter from intermediate scales onwards. Setting the unification scale at the string scale, the masses of these adjoints and the unification coupling have been calcu-




Figure 3: The range of the bottom and the top quark mass plotted against α_s . This range is obtained by varying $\tan\beta$ in the range approximately 5-60 and M_{string} in the range $2 - 7 \times 10^{17}$ GeV. Dashed lines are the prediction for the MSSM varying $\tan\beta$ in the same range.

lated. The masses of the adjoints turn out to be of the same order as that of the expected gaugino condensation scale $M_P^{2/3} m_{susy}^{1/3}$ in some cases as displayed in Figure (1) and Table 1. Taking the gauge coupling α_H for the hidden sector group G_H , to be equal to the unification coupling at the scale M_{string} , we have evolved back this value to the scale M_C , where α_H becomes non-perturbative at the two-loop order, for different choices of the hidden (matter free) group G_H . The two-loop running of the hidden sector is called for, because the scale M_C differs from the one-loop expectation (shown inside brackets in Table-1) by more than one order of magnitude. When G_H is $SU(3)$ or $SU(5)$ the scale M_C works out to be of the order of the masses of M_2 and M_3 . In comparison to MSSM, unification occurs for wider range of α_s . The bottom quark pole mass prediction from b- τ unification improves compared to that of MSSM for smaller values of α_s like 0.115. The top quark mass is evaluated assuming that the top quark Yukawa coupling at M_{string} is within domain of the quasi-infrared fixed point at the scale m_{top} . The range for the prediction of the top quark mass is consistent with experimental numbers in the moduli scenario.

$Y_t(M_{string}), Y_b(M_{string})$	$\alpha_s(m_Z)$	m_t^{pole}		m_b^{pole}		$\tan \beta$	
		<i>MSSM</i>	<i>Moduli</i>	<i>MSSM</i>	<i>Moduli</i>	<i>MSSM</i>	<i>Moduli</i>
1, 1	0.120	189	187	3.84	3.94	63.2	61.0
1, 10^{-1}	0.122	193	190	3.91	4.00	59.1	57.7
1, 10^{-2}	0.123	199	197	4.06	4.18	39.9	40.6
1, 10^{-3}	0.124	202	200	4.16	4.31	15.8	16.5
1, 10^{-4}	0.124	199	197	4.18	4.33	5.1	5.3

Table 2: The predictions of m_t^{pole} and m_b^{pole} in MSSM have been compared with those of BFY scenario for $\alpha_s(m_Z)$ which gives rise to unification in the MSSM case. For the BFY scenario $M_{string} = 5.27 \times 10^{17}$. $Y_b = h_b^2/4\pi$, where h_b is the bottom quark Yukawa coupling. The low energy value of $\tan \beta$ has been calculated using $Y_\tau(m_\tau)$ from the RGE and $m_\tau = 1.777$ GeV; and this value of $\tan \beta$ has been used to estimate the top and the bottom masses.

We thank C. Bachas and Q. Shafi for critical comments and discussions.

References

- [1] P. Ginsparg, Phys. Lett **197**, 139 (1987); V. Kaplunovsky, Nucl. Phys. **307**, 145 (1988); ibid **B382**, 436 (1992).
- [2] L. E. Ibáñez, D. Lust and G. Ross, Phys. Lett. **B272**, 251 (1991). L. E. Ibáñez and G. Ross, Nucl. Phys. **B382**, 305 (1992).
- [3] N. G. Deshpande, E. Keith and T. G. Rizzo, Phys. Rev. Lett. **70**, 3189 (1993); D. Lee and R. N. Mohapatra, Phys. Rev. **D 52**, 4125 (1995); B. Brahmachari and R. N. Mohapatra Phys. Lett. **B357**, 566 (1995); K. Benakli and G. Senjanović, ICTP report: IC/95/140, hep-ph/9507219 (to appear in Phys. Rev. **D**); K. Dienes, INS report: IASSNS-HEP-95-97, hep-th/9602045.
- [4] M. Bastero-Gil and B. Brahmachari, ICTP report-IC/96/43, hep-ph/9602391 (to appear in Nucl. Phys. **B**).

- [5] L. E. Ibáñez, Phys. Lett **B318**, 73 (1993). H. Kawabe, T. Kobayashi and N. Ohtshubo, Phys. Lett **B325**, 77 (1994). H. Kawabe, T. Kobayashi and N. Ohtshubo, Nucl. Phys. **B434**, 210 (1994); G. Aldazabal, A. Font, L. E. Ibáñez, A. M. Uranga, Nucl. Phys. **B 465**, 30 (1996).
- [6] L. Dixon, J. Harvey C. Vafa and E. Witten, Nucl. Phys. **B261**, 651 (1985); Nucl. Phys. **B274**, 285 (1986). L. Dixon, V. Kaplunovsky and J. Louis Nucl. Phys. **B355**, 649 (1991). I. Antoniadis, K. S. Narain and T. R. Taylor, Phys. Lett. **B267**, 37 (1991). S. Ferrara, C. Kounnas, D. Lust and F. Zwirner, Nucl. Phys. **B365** 431 (1991). I. Antoniadis, J. Ellis, R. Lacaze and D. V. Nanopoulos, Phys. Lett **B268**, 188 (1991).
- [7] M. K. Gaillard and R. Xiu, Phys. Lett **B2296**, 71 (1992). I. Antoniadis, J. Ellis, S. Kelly and D. V. Nanopoulos, Phys. Lett. **272B**, 31 (1991). I. Antoniadis and K. Benakli, Phys. Lett. **B295**, 219 (1992). J. Lopez, D. Nanopoulos and K. Yuan, Nucl. Phys. **B399**, 654 (1993). S. P. Martin and P. Ramond, Phys. Rev. **D51**, 6515 (1995). K. Deines and A. Faraggi, Phys. Rev. Lett. **75**, 2646 (1995).
- [8] Review of Particle Properties, Phys. Rev. **D50**, 1 (1994).
- [9] C. Bachas, C. Fabre and T. Yanagida, Phys. Lett. **B370**, 49 (1996).
- [10] H. P. Nilles, Phys. Lett. **B 115**, 193 (1982). S. Ferrara, L. Giaradello and H. P. Nilles, Phys. Lett **B125**, 457 (1983). J. P. Derendinger, L. Ibanez and H. P. Nilles, Phys. Lett **B155**, 65 (1985).
- [11] B. Pendleton and G. G. Ross, Phys. Lett. **B98**, 291 (1981); C. T. Hill, Phys. Rev. **D24**, 691 (1981). In intermediate scale models, see also, B. Brahmachari, Phys. Rev. **D52**, 1 (1995).